

Dynamics of a qubit while simultaneously monitoring its relaxation and dephasing

Q. Ficheux,^{1,2} S. Jezouin,² Z. Leghtas,^{3,2,4} and B. Huard^{1,2,*}

¹*Université Lyon, ENS de Lyon, Université Claude Bernard,
CNRS, Laboratoire de Physique, F-69342 Lyon, France*

²*Laboratoire Pierre Aigrain, Département de physique de l'ENS,
École normale supérieure, PSL Research University,*

*Université Paris Diderot, Sorbonne Paris Cité, Sorbonne Universités,
UPMC Univ. Paris 06, CNRS, 75005 Paris, France*

³*Centre Automatique et Systèmes, Mines ParisTech, PSL Research University,
60 Boulevard Saint-Michel, 75272 Paris Cedex 6, France.*

⁴*Quantic Team, INRIA Paris-Rocquencourt, Domaine de Voluceau, B.P. 105, 78153 Le Chesnay Cedex, France*

(Dated: November 6, 2017)

Decoherence originates from the leakage of quantum information into unmonitored degrees of freedom. For a qubit the two main decoherence channels are relaxation and dephasing. Here, we report an experiment on a superconducting qubit where we retrieve a significant part of the lost information in both of these channels. We demonstrate that raw averaging the corresponding measurement records provides a full quantum tomography of the qubit state where all three components of the effective spin-1/2 are simultaneously measured. From single realizations of the experiment, it is possible to infer the quantum trajectories followed by the qubit state conditioned on relaxation and/or dephasing channels. The incompatibility between these quantum measurements of the qubit leads to observable consequences in the statistics of quantum states. The high level of controllability of superconducting circuits enables us to explore many regimes from Zeno effect to underdamped Rabi oscillations depending on the relative strengths of driving, dephasing and relaxation.

Decoherence can be understood as the result of unread measurements of a system by its environment. For a qubit, the two main sources of decoherence are relaxation by spontaneous emission and dephasing by the unwanted readout of coupled quantum systems (Fig. 1a). What becomes of the qubit state if, instead of disregarding the information leaking to the environment, we continuously monitor both decoherence channels? Owing to measurement backaction, the knowledge of the measurement record then leads to a stochastic quantum trajectory of the qubit state for each single realization of an experiment [1–3]. Recently, diffusive quantum trajectories were observed following the continuous homodyne or heterodyne measurements of either a dephasing channel [4–9] or a relaxation channel [10, 11]. Here we report an experiment in which we have simultaneously monitored the spontaneous emission of a superconducting qubit by heterodyne measurement (relaxation channel) and the transmitted field through a dispersively coupled cavity by homodyne measurement (dephasing channel). We demonstrate that the average outcomes of these two non-projective measurements are the three coordinates x , y and z of the Bloch vector. It is remarkable that a full quantum tomography can be obtained by simply raw averaging measurement outcomes of many realizations of a single experiment despite the incompatibility of the three observables that characterize a qubit state. For single realizations the resulting quantum trajectories show signatures of the incompatibility between the measurement channels, therefore extending the previously explored case of two incompatible measurement outcomes [10, 12] to the case of three spin directions.

By varying the drive amplitudes at the cavity and qubit transition frequencies, we are able to reach a variety of regimes corresponding to different configurations for Ω/Γ_1 and Γ_d/Γ_1 , where Ω is the Rabi frequency, Γ_1 the fixed relaxation rate and Γ_d the dephasing rate. This work hence provides a textbook experimental demonstration of quantum measurement backaction on a qubit with incompatible and simultaneous measurements.

Two parallel detection setups operate via spatially separated measurement lines (see Fig. 1b). The fluorescence heterodyne detection setup enables the measurement of both quadratures $u(t)$ and $v(t)$ of the spontaneously emitted field out of a 3D transmon qubit [13]. The complex amplitude of the emitted field is on average proportional to the expectation of the qubit lowering operator $\sigma_- = (\sigma_x - i\sigma_y)/2$ so that u and v are on average proportional to the expectations of σ_x and σ_y . Here, $\sigma_x = |g\rangle\langle e| + |e\rangle\langle g|$, $\sigma_y = i|g\rangle\langle e| - i|e\rangle\langle g|$ and $\sigma_z = |e\rangle\langle e| - |g\rangle\langle g|$ are the qubit Pauli operators, where $|g\rangle$ and $|e\rangle$ are the ground and excited states. For a single realization, the measurement outcomes read [10]

$$\begin{cases} u(t)dt = \sqrt{\eta_f\Gamma_1/2}x(t)dt + dW_u(t) \\ v(t)dt = \sqrt{\eta_f\Gamma_1/2}y(t)dt + dW_v(t) \end{cases}, \quad (1)$$

where $\Gamma_1 = (15 \mu\text{s})^{-1}$ is the qubit relaxation rate, $\eta_f = 0.14$ is the total fluorescence measurement efficiency, $x(t) = \text{Tr}(\sigma_x\rho_t)$ and $y(t) = \text{Tr}(\sigma_y\rho_t)$ are the qubit Bloch coordinates corresponding to the density matrix ρ_t (see Fig. 1a) and W_u and W_v are two independent stochastic Wiener processes describing the measurement noise, which includes the zero point fluctuations, and such that $dW^2 = dt$ and dW is zero on average. Ex-




FIG. 1: Scheme of the measurement setup and quantum trajectory resulting from its three outputs. **a.** Bloch vector representation of a qubit whose state is described by a density matrix $\rho_t = (\mathbf{1} + x(t)\sigma_x + y(t)\sigma_y + z(t)\sigma_z)/2$. A quantum trajectory ρ_t is represented as a blue line. The qubit decoherence can be modeled as originating from a relaxation channel at a rate Γ_1 and a dispersive measurement channel at a rate Γ_d . **b.** A superconducting qubit in a cavity is driven by two microwave signals at the weakly coupled input. The one at qubit frequency $f_q = 5.353$ GHz (orange) induces Rabi oscillations of the qubit at frequency Ω . The one at cavity frequency $f_d = 7.761$ GHz (purple) leads to a dispersive measurement of the qubit state along σ_z . A diplexer at the strongly coupled output port separates the outgoing signals depending on their frequency. The radiation at f_q that is spontaneously emitted by the qubit is processed by a Josephson Parametric Converter (JPC) [14, 15] so that a following heterodyne measurement reveals the two quadratures $u(t)$ and $v(t)$ of the fluorescence field [10, 16, 17]. The transmitted signal at f_d is processed by a doubly pumped Josephson Parametric Amplifier (JPA) [4, 18] with a pump phase such that a following homodyne measurement reveals the quadrature $w(t)$ of the field at f_d . **c.** Measurement records u, v and w as a function of time for one realization of the experiment. These records feed the stochastic master equation (3), which leads to the trajectory in **a.**

perimentally, the measurement takes a non infinitesimal time $dt = 100$ ns, which we chose smaller than the inverse measurement rates and compatible with the detection bandwidth [19].

Similarly, the dispersive detection setup (see Fig. 1b) enables the measurement of a single quadrature $w(t)$ of the transmitted field at frequency $f_d = f_r - \chi_{cq}/2$, which is between the cavity resonance frequencies f_r and $f_r - \chi_{cq}$ respectively corresponding to a qubit in the ground and excited state ($\chi_{cq} = 2\pi \times 5.1$ MHz). The phase of the measured quadrature in the homodyne measurement can then be chosen in such a way that [8, 17]

$$w(t)dt = \sqrt{2\eta_d\Gamma_d}z(t)dt + dW_w(t). \quad (2)$$

Importantly, the measurement induced dephasing rate Γ_d can be tuned arbitrarily as it is proportional to the drive power at f_d . Similarly to the notations above, $\eta_d = 0.34$ is the total dispersive measurement efficiency, $z(t) = \text{Tr}(\sigma_z\rho_t)$ is the last of the three Bloch coordinates (see Fig. 1a) and W_w is another independent Wiener process.

As can be seen from Eqs. (1,2), taking a raw average of the outcomes (u, v, w) on a large number of realizations of the experiment directly leads to the Bloch coordinates (x, y, z) of the qubit. In Fig. 2, we show the direct averaging of the three outcomes in two configurations of the input drives: one in the regime of underdamped Rabi oscillations (Fig. 2a) and another in the regime of strong dispersive measurement rate, the so-called Zeno regime (Fig. 2b). The raw averaging of (u, v, w) , once rescaled by the prefactors in Eqs. (1,2), agrees well with the average evolution of the qubit, as predicted by the solution of the master equation (Eq. 3 below without the last stochastic term). We thus demonstrate that performing a dispersive measurement and a measurement of fluorescence reveals information on all three components of a spin-1/2.




FIG. 2: Rescaled averaging of the three measurement records directly gives the average evolution of the qubit state in Bloch representation. **a.** Dots: Rescaled average of the measurement records $\tilde{u}(t) = \bar{u}(t)/\sqrt{\eta_f\Gamma_1/2}$, $\tilde{v}(t) = \bar{v}(t)/\sqrt{\eta_f\Gamma_1/2}$ and $\tilde{w}(t) = \bar{w}(t)/\sqrt{2\eta_d\Gamma_d}$ for $1.5 \cdot 10^6$ realizations of an experiment where the qubit starts in $|g\rangle$ at time 0 and is driven so that it rotates at a Rabi frequency $\Omega/2\pi = (2 \mu\text{s})^{-1}$ around σ_y and endures a measurement induced dephasing rate $\Gamma_d = (5 \mu\text{s})^{-1}$. Lines: Calculated coordinates of the Bloch vector $x(t)$, $y(t)$ and $z(t)$ from the master equation (Eq. 3 with $\eta_i = 0$). **b.** Same figure in the Zeno regime with a drive such that $\Omega/2\pi = (16 \mu\text{s})^{-1}$ and $\Gamma_d = (0.9 \mu\text{s})^{-1}$.

Experiments of Figs. 2a and 2b differ by the relative rate of the dispersive readout Γ_d compared to the Rabi frequency Ω . For weak measurement rate $\Gamma_d, \Gamma_1 < \Omega$, the Rabi oscillations are underdamped, while they are overdamped when $\Gamma_d \gg \Omega, \Gamma_1$ owing to the fact that the Zeno effect prevents any unitary evolution such as Rabi oscillations. For a single realization of the experiment though, the trajectory of the qubit state that one can infer from the measurement records $u(t)$, $v(t)$ and $w(t)$




FIG. 3: Correlations between the coordinates $(x_{\text{traj}}, y_{\text{traj}}, z_{\text{traj}})$ of the trajectories after $19.8 \mu\text{s}$ of evolution and an independent tomography on the dataset corresponding to the experiment of Fig. 2a. Average value of the tomography results for the subset of trajectories ending up less than 0.01 away from a given value of x_{traj} (Fig. a), y_{traj} (Fig. b) or z_{traj} (Fig. c). The error bars are given by the standard deviation of the tomography results divided by the square root of the number of trajectories in the subset. The agreement between the tomography and the coordinates of the trajectories demonstrates the validity of the quantum trajectories. Fig. d. Bloch sphere representation of 3 quantum trajectories that end up with $0.74 < x_{\text{traj}} < 0.76$ (red dashed line) after $19.8 \mu\text{s}$ corresponding to one bin of the histogram in Fig. a.

can strongly differ from this average behavior. In order to determine this quantum trajectory, one can use the formalism of the stochastic master equation [17]. The density matrix at time $t + dt$ can be decomposed into $\rho_{t+dt} = \rho_t + d\rho_t$, where

$$d\rho_t = i\left[\frac{\Omega}{2}\sigma_y, \rho_t\right]dt + \sum_k D_k(\rho_t)dt + \sum_k \sqrt{\eta_k} \mathcal{M}_k(\rho_t) dW_k(t), \quad (3)$$

with the four Lindblad superoperator ($k \in \{u, v, w, \varphi\}$)

$$\mathcal{D}_k(\rho_t) = L_k \rho_t L_k^\dagger - \frac{1}{2} \rho_t L_k^\dagger L_k - \frac{1}{2} L_k^\dagger L_k \rho_t, \quad (4)$$

and the measurement backaction superoperator

$$\mathcal{M}_k(\rho_t) = L_k \rho_t + \rho_t L_k^\dagger - \text{Tr}(L_k \rho_t + \rho_t L_k^\dagger) \rho_t. \quad (5)$$

In these expressions, the jump operators corresponding to heterodyning fluorescence are $L_u = \sqrt{\Gamma_1/2}\sigma_-$ and $L_v = i\sqrt{\Gamma_1/2}\sigma_-$ and the jump operator corresponding to homodyning the dispersive measurement is $L_w = \sqrt{\Gamma_d/2}\sigma_z$. A fourth jump operator $L_\varphi = \sqrt{\Gamma_\varphi/2}\sigma_z$ corresponds to the unread ($\eta_\varphi = 0$) pure dephasing of the qubit, so that the total decoherence rate $\Gamma_2 = \frac{\Gamma_1}{2} + \Gamma_\varphi + \Gamma_d$ can be tuned from $\Gamma_2 = (11.2 \mu\text{s})^{-1}$ to higher arbitrary values depending on the power of the drive at frequency f_d .

Using this formalism it is possible to reconstruct the quantum trajectory of the qubit state in time from any set of measurement records (see Fig. 1a,c in the case where $\Omega/2\pi = (5.2 \mu\text{s})^{-1}$ and $\Gamma_d = (0.9 \mu\text{s})^{-1}$). The validity of the reconstructed quantum trajectories can be tested independently by post-selecting an ensemble of realizations of the experiment for which the trajectory predicts a given value $x(T) = x_{\text{traj}}$ at a time T . If the trajectories are valid, then a strong measurement of σ_x at time T should give x_{traj} on average on this post-selected ensemble of realizations (Fig. 3a). We have checked for any value of x_{traj} , y_{traj} and z_{traj} (Fig. 3), and for 30 representative configurations of drives that the trajectories predict the strong measurement results [19]. In fact, we found that the agreement is verified for efficiencies η_f and η_d within a confidence interval of ± 0.02 for any of the 30 configurations.

Any measurement record is a stochastic process and the corresponding quantum trajectories follow a random walk in the Bloch sphere with a state dependent diffusion constant. The inherent backaction of a quantum measurement is thus better discussed by representing distributions of states at a given time [4, 5, 8, 10–12, 20] or distributions of trajectories for a given duration [7, 21–24]. Figure 4 gives a different perspective to the Rabi oscillation of Fig. 2a by representing the distributions of the qubit states conditioned on the three measurement records $u(t)$, $v(t)$ and $w(t)$ for 1.5 million realizations of the experiment. In the supplemental material [19], one can find movies of the distributions of 1.5 millions experimental realizations for each configuration of the Rabi frequency Ω and the dephasing rate Γ_d for a set of 30 different experimentally realized configurations. Evidently, the Rabi drive term $-\Omega\sigma_y/2$ still provides an overall angular velocity in the $x-z$ plane of the Bloch sphere. However, the measurement backaction is such that some trajectories are delayed while others are advanced compared to the average evolution. As time increases the spread in the qubit states grows as a result of the cumulated effect of the stochastic measurement backaction at each time step.

The effect of decoherence under a strong Rabi drive corresponds to an average loss of purity, defined as $\text{Tr}(\rho^2) = (1 + x^2 + y^2 + z^2)/2$ and it can be seen as a decreasing distance of the mean trajectory from the center of the Bloch sphere when time increases (solid line). When the dispersive measurement (dephasing channel) is measured in presence of the Rabi drive around σ_y , the corresponding distribution of states tends to be uniform in the $x-z$ plane at long times (right panel in Fig. 4b), which is similar to what is obtained by simultaneously measuring σ_x and σ_z in an effectively undriven qubit [12]. The experiment thus illustrates the fact that the average loss of purity corresponds to the statistical uncertainty on the quantum state when the decoherence channel is unread.




FIG. 4: Evolution of the distribution of quantum states in the same conditions as Fig. 2a. Colored dots: Each frame represents the marginal distribution, in the $x - y$ (Fig. a), $x - z$ (Fig. b) and $y - z$ (Fig. c) planes of the Bloch sphere, of the states of the qubit at a given time τ for 1.5 million realizations of the experiment. Each state (x, y, z) is reconstructed from the measurement records $\{u(t), v(t), w(t)\}$ from time t between 0 to τ using Eq. (3). Time τ is increasing from $0.3 \mu\text{s}$ to $19 \mu\text{s}$ from left to right as indicated at the bottom of the figure. For each figure, the surrounding black circles represent the pure states of the plane (e.g. $z = 0$ for Fig. a). Solid lines: average projection of all 1.5 millions of quantum trajectories $\{x(t), y(t), z(t)\}$ for $0.2 \mu\text{s} < t < \tau$.

Interestingly, while the average trajectory stays in the $x - z$ plane with $\langle \sigma_y \rangle = 0$, the backaction of the fluorescence measurement leads to a nonzero spread in the y direction of the Bloch sphere. This competition between the backaction of relaxation (fluorescence measurement) and dephasing (dispersive measurement) measurements can be better observed when decoherence dominates the dynamics. In Fig. 5, we show the distributions of qubit states at a long time $\tau = 6.5 \mu\text{s}$ after which the distribution is close to its steady state while the qubit is both Rabi driven and dispersively measured at a strong measurement rate. The trajectories are determined using three sets of measurement records: dispersive only $\{w(t)\}$, fluorescence only $\{u(t), v(t)\}$ or both. As in Fig. 2b, the Zeno effect then leads to the dampening of the Rabi oscillations and the average trajectory (solid line) quickly reaches its steady state.

In contrast, a trajectory corresponding to a single realization of the experiment where the dispersive measurement $w(t)$ is recorded is found to consist in a series of stochastic jumps between two areas of the Bloch sphere that are close to the two eigenstate of the σ_Z measurement operator. In the distribution of states, this leads to two areas with high probability of occupation near the poles of the Bloch sphere. These areas can be interpreted as zones frozen by the Zeno effect. The rest of the Bloch sphere is still occupied with a lower probability

(Fig. 5b) because of the finite time it takes for the jump to occur from one pole to the next under strong dispersive measurement rate [25]. Note how the ensemble of trajectories can go from uniform for weak measurement rates (Fig. 4b rightest panel) to localized at the poles for strong measurement rates (Fig. 5b).

As can be seen from Figs. 5a,c, the dispersive measurement alone does not provide any backaction towards the y direction of the Bloch sphere so that the qubit states keeps a zero σ_y component during its evolution. This is in stark contrast with the trajectories corresponding to measurement records $\{u(t), v(t)\}$ of the fluorescence (Figs. 5d-f), where at long times the qubit states spans a small ball in the Bloch sphere. Therefore, the combined action of Rabi drive and fluorescence measurement backaction leads to a uniform spread of the qubit state close to the most entropic state $\mathbf{1}/2$ at the center of the sphere. As expected, the quantum states that are conditioned on all measurement records $\{u(t), v(t), w(t)\}$ are less entropic than with a single measurement. This can be seen in Figs. 5g-i where the spread of the distributions is larger than for the cases of single measurements Figs. 5a-f.

A clear asymmetry appears in the spread of the marginal distribution in the $x - y$ plane of Fig. 5g between positive and negative values of x . This asymmetry originates from the fact that the fluorescence mea-




FIG. 5: Impact of the type of detector on the distribution of quantum states after $\tau = 6.5 \mu\text{s}$ in the Zeno regime ($\Omega/2\pi = (5.2 \mu\text{s})^{-1} \ll \Gamma_d = (0.9 \mu\text{s})^{-1}$). **a, b, c.** Marginal distribution in the x, y (Fig. **a**), $x - z$ (Fig. **b**) and $y - z$ (Fig. **c**) planes of the Bloch sphere of the qubit states ρ_τ corresponding to 1.5 millions of measurement records at the cavity frequency only $\{w(t)\}$ from time t between 0 and τ . As in Fig. 4, the boundary of the Bloch sphere is represented as a black circle and the average quantum trajectory as a solid line. **d. e. f.** Case where the states are conditioned on fluorescence records $\{u(t), v(t)\}$ instead. **g. h. i.** Case where the states are conditioned on both fluorescence and dispersive measurement records $\{u(t), v(t), w(t)\}$.

surement is linked to the jump operator σ_- for which $|g\rangle$ is the single pointer state. Indeed the measurement backaction is null when the qubit state is close to $|g\rangle$ ($\mathcal{M}_u(|g\rangle\langle g|) = \mathcal{M}_v(|g\rangle\langle g|) = 0$) while it is strongest when the qubit state is close to $|e\rangle$. Since the Rabi drive correlates the ground state to positive x (red zone shifted to the right of the south pole in Fig. 5h) and the excited state to negative x , the spread in y is smaller for positive x than for negative x . This asymmetry highlights the profound difference between measuring both quadratures of fluorescence and measuring σ_x and σ_y simultaneously using dynamical states as in Ref. [12, 26]. While both methods lead to the same result on average, their backaction differs. The latter corresponds to quantum non-demolition measurements, while fluorescence does not. In the end, the asymmetry in the distributions of Figs. 5g,i results from the incompatibility between a dis-

persive measurement with no backaction on $|e\rangle$ and a fluorescence measurement with maximal backaction on $|e\rangle$.

In conclusion, we have shown quantum trajectories of a superconducting qubit reconstructed from three measurements originating from the simultaneous monitoring of its decoherence channels. It looks promising to test new statistical properties of quantum trajectories [27, 28], fluctuation relations in quantum thermodynamics [29–35], quantum smoothing protocols [20, 36–41], and to perform parameter estimation [42, 43].

Acknowledgements We thank Philippe Campagne-Ibarcq, Michel Devoret, Andrew Jordan, Raphaël Lescanne, Mazyar Mirrahimi, Klaus Mølmer, Pierre Rouchon, Alain Sarlette, Irfan Siddiqi and Pierre Six for fruitful interactions. Nanofabrication has been made within the consortium Salle Blanche Paris Centre. This work was supported by the EMERGENCES grant QUMOTEL of Ville de Paris. Q.F. and S.J. contributed equally to this work.

* Electronic address: benjamin.huard@ens-lyon.fr

- [1] H. Carmichael, *An Open Systems Approach to Quantum Optics* (Springer Berlin Heidelberg, 1993).
- [2] A. Barchielli and M. Gregoratti, *Quantum Trajectories and Measurements in Continuous Time*, vol. 782 (Springer-Verlag Berlin Heidelberg, 2009).
- [3] H. M. Wiseman and G. J. Milburn, *Quantum Measurement and Control* (Cambridge University Press, 2009).
- [4] K. W. Murch, S. J. Weber, C. Macklin, and I. Siddiqi, *Nature* **502**, 211 (2013).
- [5] M. Hatridge, S. Shankar, Mirrahimi, F. Schackert, K. Geerlings, T. Brecht, K. Sliwa, B. Abdo, L. Frunzio, Girvin, et al., *Science* **339**, 178 (2013).
- [6] G. de Lange, D. Ristè, M. Tiggelman, C. Eichler, L. Tornberg, G. Johansson, a. Wallraff, R. Schouten, and L. DiCarlo, *Physical Review Letters* **112**, 080501 (2014).
- [7] S. J. Weber, a. Chantasri, J. Dressel, a. N. Jordan, K. W. Murch, and I. Siddiqi, *Nature* **511**, 570 (2014).
- [8] S. J. Weber, K. W. Murch, M. E. Kimchi-Schwartz, N. Roch, and I. Siddiqi, *Comptes Rendus Physique* **17**, 766 (2016).
- [9] A. Chantasri, M. E. Kimchi-Schwartz, N. Roch, I. Siddiqi, and A. N. Jordan, *Physical Review X* **6**, 041052 (2016).
- [10] P. Campagne-Ibarcq, P. Six, L. Bretheau, A. Sarlette, M. Mirrahimi, P. Rouchon, and B. Huard, *Phys. Rev. X* **6**, 11002 (2016).
- [11] M. Naghiloo, N. Foroozani, D. Tan, A. Jadbabaie, and K. W. Murch, *Nature Communications* **7**, 11527 (2016).
- [12] S. Hacohe-Gourgy, L. S. Martin, E. Flurin, V. V. Ramasesh, K. B. Whaley, and I. Siddiqi, *Nature* **538**, 491 (2016).
- [13] H. Paik, D. I. Schuster, L. S. Bishop, G. Kirchmair, G. Catelani, a. P. Sears, B. R. Johnson, M. J. Reagor, L. Frunzio, L. I. Glazman, et al., *Physical Review Letters* **107**, 240501 (2011).
- [14] N. Bergeal, F. Schackert, M. Metcalfe, R. Vijay, V. E. Manucharyan, L. Frunzio, D. E. Prober, R. J. Schoelkopf,

- S. M. Girvin, and M. H. Devoret, *Nature* **465**, 64 (2010).
- [15] N. Roch, E. Flurin, F. Nguyen, P. Morfin, P. Campagne-Ibarcq, M. H. Devoret, and B. Huard, *Physical Review Letters* **108**, 147701 (2012).
- [16] P. Campagne-Ibarcq, L. Bretheau, E. Flurin, A. Auffèves, F. Mallet, and B. Huard, *Phys. Rev. Lett.* **112**, 180402 (2014).
- [17] P. Campagne-Ibarcq, Ph.D. thesis, Ecole Normale Supérieure (ENS) (2015).
- [18] A. Kamal, A. Marblestone, and M. Devoret, *Physical Review B* **79**, 184301 (2009).
- [19] Supplemental Material (2017).
- [20] D. Tan, S. Weber, I. Siddiqi, K. Mølmer, and K. Murch, *Physical Review Letters* **114**, 090403 (2015).
- [21] A. N. Jordan, A. Chantasri, P. Rouchon, and B. Huard, *Quantum Studies: Mathematics and Foundations* **3**, 237 (2016).
- [22] M. Naghiloo, D. Tan, P. M. Harrington, P. Lewalle, A. N. Jordan, and K. W. Murch, arxiv:1612.03189 (2016).
- [23] A. Chantasri, M. E. Kimchi-Schwartz, N. Roch, I. Siddiqi, and A. N. Jordan, *Physical Review X* **6**, 041052 (2016).
- [24] P. Lewalle, A. Chantasri, and A. N. Jordan, *Physical Review A* **95**, 042126 (2017).
- [25] Y. Choi and A. N. Jordan, *Physical Review A* **88**, 052128 (2013).
- [26] U. Vool, S. Shankar, S. O. Mundhada, N. Ofek, A. Narla, K. Sliwa, E. Zalys-Geller, Y. Liu, L. Frunzio, R. J. Schoelkopf, et al., *Physical Review Letters* **117**, 133601 (2016).
- [27] A. Franquet and Y. V. Nazarov, *Physical Review B* **95**, 085427 (2017).
- [28] A. Chantasri, J. Atalaya, S. Hacothen-Gourgy, L. S. Martin, I. Siddiqi, and A. N. Jordan, arxiv:1706.09670 (2017).
- [29] Z. Gong, Y. Ashida, and M. Ueda, *Physical Review A* **94**, 012107 (2016).
- [30] C. Elouard, D. A. Herrera-Martí, M. Clusel, and A. Auffèves, *npj Quantum Information* **3**, 9 (2017).
- [31] J. J. Alonso, E. Lutz, and A. Romito, *Physical Review Letters* **116**, 080403 (2016).
- [32] P. G. Di Stefano, J. J. Alonso, E. Lutz, G. Falci, and M. Paternostro, arxiv:1704.00574 (2017).
- [33] S. Deffner, J. P. Paz, and W. H. Zurek, *Physical Review E* **94**, 010103 (2016).
- [34] M. Naghiloo, D. Tan, P. M. Harrington, J. J. Alonso, E. Lutz, A. Romito, and K. W. Murch, arxiv:1703.05885 (2017).
- [35] F. Liu, arxiv:1710.02311 (2017).
- [36] M. Tsang, *Physical Review Letters* **102**, 250403 (2009).
- [37] M. Tsang, *Physical Review A* **80**, 033840 (2009).
- [38] S. Gammelmark, B. Julsgaard, and K. Mølmer, *Physical Review Letters* **111**, 160401 (2013).
- [39] I. Guevara and H. Wiseman, *Physical Review Letters* **115**, 180407 (2015).
- [40] D. Tan, M. Naghiloo, K. Mølmer, and K. W. Murch (2016).
- [41] D. Tan, N. Foroozani, M. Naghiloo, A. H. Küllerich, K. Mølmer, and K. W. Murch (2017).
- [42] P. Six, P. Campagne-Ibarcq, L. Bretheau, B. Huard, and P. Rouchon, in *Decision and Control (CDC), 2015 IEEE 54th Annual Conference on (IEEE, 2015)*, pp. 7742–7748.
- [43] A. H. Küllerich and K. Mølmer, *Physical Review A* **94**, 032103 (2016).